

NUS/HEP/920503

May 1992

Duality in Multi-layered Quantum Hall Systems

CHRISTOPHER TING

Defence Science Organization, 20 Science Park Drive, Singapore 0511

Abstract

The braid group dynamics captures the fractional quantum Hall effect (FQHE) as a manifestation of puncture phase. When the dynamics is generalized for particles on a multi-sheeted surface, we obtain new tools which determine the fractional charges, the quantum statistics, and the filling factors of the multi-layered FQHE. A many-quasi-hole wavefunction is proposed for the bilayered samples. We also predict a $\nu = 5/7$ FQHE for triple-layered samples. The viability of *3-dimensional* FQHE and the application of the concept of generalized duality to anyonic superconductivity are discussed.

PACS numbers: 71.28, 71.10, 72.20M

Laughlin's theory [1] of fractional quantum Hall effect (FQHE) elegantly describes the incompressibility of the 2-dimensional electronic system under the influence of a strong, uniform magnetic field. The theory is pivoted on an ansatz which is taken to be the ground state ψ_m of the many-body system displaying FQHE:

$$\psi_m = \prod_{a < b} (w_a - w_b)^m \exp\left(-\frac{1}{4\ell^2} \sum_a |w_a|^2\right), \quad (1)$$

where ℓ denotes the magnetic length and w_a are the coordinates of N electrons. Now, this ansatz is for single-layered quantum Hall system, with all the spins of the electrons polarized in the direction of the magnetic field. Since electrons are fermions, in order to have ψ_m anti-symmetric with respect to the permutation of indices a , m must be an odd number. Recently, however, two experimental groups [2, 3] successfully produced a $\nu = \frac{1}{2}$ FQHE. The samples they used were double-layered 2-dimensional electronic systems. Labelling the two layers and treating them as additional degrees of freedom, the $\nu = \frac{1}{2}$ effect was predicted by Yoshioka, MacDonald and Girven (YMG) [4]. Basically, they made a straightforward generalization of Laughlin's ansatz (1) as follows:

$$\begin{aligned} \psi_{m_1, m_2, n}(w_a^{[1]}, w_a^{[2]}) &= \prod_{a < b}^{N^{[1]}} (w_a^{[1]} - w_b^{[1]})^{m_1} \prod_{a < b}^{N^{[2]}} (w_a^{[2]} - w_b^{[2]})^{m_2} \prod_{a, b}^{N^{[1]}, N^{[2]}} (w_a^{[1]} - w_b^{[2]})^n \\ &\times \exp -\frac{1}{4\ell^2} \left(\sum_a^{N^{[1]}} |w_a^{[1]}|^2 + \sum_a^{N^{[2]}} |w_a^{[2]}|^2 \right). \end{aligned} \quad (2)$$

Presumably, the two labels [1] and [2] are some analogue of spin, as $\psi_{m_1, m_2, n}$ was first discussed by Halperin [5] in the context of single-layered *unpolarized* quantum Hall systems. On the other hand, the novel thing about $\psi_{m_1, m_2, n}$ in YMG's analysis is to interpret $\prod_{a, b} (w_a^{[1]} - w_b^{[2]})^n$ as inter-layer correlation. They concluded that $\psi_{m_1, m_2, n}$ could well be a good ansatz for the ground states of doubled-layered FQHE *if* the distance d between the layers is about 1.4 times of ℓ for $\psi_{3,3,1}$ and 3ℓ for $\psi_{5,5,1}$. To reveal the underlying physics of $\psi_{m_1, m_2, n}$, it is important to find a Hamiltonian for which $\psi_{m_1, m_2, n}$ is an *exact* ground state and to study the nature of the quasi-excitations of the double-layered FQHE. What is the value of the fractional charge carried by each quasi-hole? What is the quantum statistics of the quasi-holes? Is it possible to have a theoretical tool that can generalize the analysis of these issues even to *multi*-layered FQHE? These are the questions we are going to answer.

Earlier, we have focussed on the *topological* underpinning of the single-layered FQHE [6, 7] from the braid group approach [8]. It was understood that Laughlin wavefunction ψ_m satisfied the ground state equations of a Hamiltonian suggested by the representation theory of the Artin braid group. The braid group approach reveals that FQHE

is a manifestation of the puncture phase, wherein each electron sees its counterparts as punctures. When the 2-dimensional systems enter the puncture phase, the configuration space for each particle is no longer simply connected. The non-trivial topology necessitates the consideration of homotopic constraint that each possible path of a particle is imposed upon when travelling from one location to the other. In this regard, Y. S. Wu explicitly shows that the 1-dimensional path integral representation of Artin braid group leads to a generalized quantum statistics interpolating between bosons and fermions [9]. We introduced the concept of *charged* winding “number” and thereby expanded Wu’s construction to include the possibility of non-abelian anyons [8, 10]. Beside being a non-abelian generalization of Wu’s representation theory, our formulation of homotopical constraint with the charged winding number directly points out the relevance of Knizhnik-Zamolodchikov equations [11] to the physics of anyons. Indeed, it can be readily seen that Laughlin’s wavefunction is an *exact* solution of

$$(D_a + \frac{B}{4}\bar{w}_a)\psi_m = 0, \quad (3)$$

where

$$D_a \equiv \frac{\partial}{\partial w_a} - m \sum_{b=1, b \neq a}^N \frac{1}{w_a - w_b} \quad (4)$$

is the Knizhnik-Zamolodchikov operator for the conformal blocks of the current algebra. We are working in a unit system where all the universal constants are equal to 1 and B is the magnetic field strength. Using the same approach, we have also captured $\psi_{m_1, m_2, n}$ as an *exact* solution of

$$\left(\frac{\partial}{\partial w_a^{[1]}} - m_1 \sum_{b \neq a}^{N^{[1]}} \frac{1}{w_a^{[1]} - w_b^{[1]}} - n \sum_b^{N^{[2]}} \frac{1}{w_a^{[1]} - w_b^{[2]}} + \frac{B}{4}\bar{w}_a^{[1]} \right) \psi_{m_1, m_2, n} = 0. \quad (5)$$

$$\left(\frac{\partial}{\partial w_a^{[2]}} - n \sum_b^{N^{[1]}} \frac{1}{w_a^{[2]} - w_b^{[1]}} - m_2 \sum_{b \neq a}^{N^{[2]}} \frac{1}{w_a^{[2]} - w_b^{[2]}} + \frac{B}{4}\bar{w}_a^{[2]} \right) \psi_{m_1, m_2, n} = 0. \quad (6)$$

This may appear as a trivial generalization of (3). The physics behind, however, reaches much deeper than that [12]. It is shown that (5, 6) are the ground state equations of particles living in the *double-sheeted* surface. The strength n with which particles from the first sheet see those in the second sheet as punctures can now differ from m_1 , and vice versa. As a consequence, the possibility of *mutual* statistics between the particles of different layers arises, as originally discussed by Wilczek [13] in the language of adiabatic transport. In order for $\psi_{m_1, m_2, n}$ to be an anti-symmetric wavefunction, m_1, m_2 and n are odd integers. This is because we are describing a system of electrons, which are fermions. We realize that the case with $n = m_1 = m_2$ is trivial. This is simply the

redundant way of saying that the two layers are effectively indistinguishable and the two equations (5, 6) collapse back to (3). Physically, it corresponds to the situation where the distance d between the layers is zero. Therefore, for the Hamiltonian suggested by the representation theory of braid group over the double-sheeted surface to be a truly non-trivial generalization, the pathological case is ruled out. Next, one would expect n to be no larger than m_1 and m_2 . This is because the inter-layer correlation should be weaker than the intra-layer correlation. The strength m_i with which particles of layer i see those of the same layer as punctures should be larger than n with which to see those of other layer. Let us summarize the physical arguments: (A) m_1, m_2, n are odd integers, (B) $n = m_1 = m_2$ is forbidden, and (C) $n \leq \min(m_1, m_2)$. A succinct way to express the relevant quantum numbers of the double-layered system is to write down a matrix of topological coupling strengths after (5, 6):

$$\begin{pmatrix} m_1 & n \\ n & m_2 \end{pmatrix} \quad (7)$$

Curiously, the physical inputs (A), (B) and (C) imply a condition to be satisfied by (7): $m_1 m_2 - n^2 > 0$. Therefore, the coupling matrix for $\psi_{m_1, m_2, n}$ is non-singular.

By looking at $\psi_{m_1, m_2, n}$ alone, it is not clear how the unusual quantum statistics may arise. After all, since m_1, m_2, n are odd integers, the quantum statistics of the two layers of electrons characterized by $\psi_{m_1, m_2, n}$ is still the usual one we know from the Pauli principle. To appreciate the meaning of mutual statistics, we have to consider the quasi-excitations. To warm up, let us briefly sketch how the quasi-hole wavefunctions of the single-layered systems can be obtained from the braid group dynamics. Suppose the charge of each quasi-hole is c , and we use $u_a, a = 1, \dots, N_h$ to denote their coordinates. It turns out that the many-quasi-hole state ψ_m^{qh} of ψ_m satisfies the following equations:

$$\begin{aligned} \left(\partial_{w_a} + \frac{B}{4} \bar{w}_a - m \sum_{b=1, b \neq a}^N \frac{1 \times 1}{w_a - w_b} - m \sum_{b=1}^{N_h} \frac{1 \times c}{w_a - u_b} \right) \psi_m^{qh} &= 0, \\ \left(\partial_{u_a} + \frac{cB}{4} \bar{u}_a - m \sum_{b=1}^N \frac{c \times 1}{u_a - w_b} - m \sum_{b=1, b \neq a}^{N_h} \frac{c \times c}{u_a - u_b} \right) \psi_m^{qh} &= 0. \end{aligned} \quad (8)$$

One readily solves the equations and obtains

$$\psi_m^{qh}(u_1, \dots, u_{N_h}; w_1, \dots, w_N) = \prod_{1 \leq a < b \leq N_h} (u_a - u_b)^c \exp\left(-\frac{c}{4\ell^2} \sum_a |u_a|^2\right) \prod_{a,b} (u_a - w_b) \psi_m \quad (9)$$

It is well known that the quasi-holes are fractionally charged entities. We shall see later how the value of c is determined. As can be seen from (9), the statistics of the quasi-holes is $\theta/\pi = c$. Observe that there exists a kind of “duality” in (8). Namely, if

we rewrite it in the following form

$$\begin{aligned} \left(\partial_{w_a} + \frac{B}{4} \bar{w}_a - m \sum_{b=1, b \neq a}^N \frac{1 \times 1}{w_a - w_b} - mc \sum_{b=1}^{N_h} \frac{1 \times 1}{w_a - u_b} \right) \psi_m^{qh} &= 0, \\ \left(\partial_{u_a} + \frac{cB}{4} \bar{u}_a - mc \sum_{b=1}^N \frac{1 \times 1}{u_a - w_b} - mc^2 \sum_{b=1, b \neq a}^{N_h} \frac{1 \times 1}{u_a - u_b} \right) \psi_m^{qh} &= 0. \end{aligned} \quad (10)$$

we have a striking resemblance with the double-layered systems (5, 6). In words, the quasi-holes of charge c can be regarded as electrons (of unit charge) living in the second layer! Viewed in this light, the coupling matrix of such fictitious double-layered system is

$$\begin{array}{cc} & \begin{matrix} e & qh \end{matrix} \\ \begin{matrix} e \\ qh \end{matrix} & \left(\begin{array}{cc} m & mc \\ mc & mc^2 \end{array} \right) \end{array} \quad (11)$$

The crucial difference between two layers of real electrons and one layer of electrons and quasi-holes is that while the coupling matrix of the former is non-singular, that of the latter is singular. That the coupling matrix for electron-quasi-hole system (11) is singular is a manifestation of the electric-magnetic duality. The charges c of the quasi-holes are related to the number of extra flux units being buffered by the Laughlin fluid when the background magnetic field strength B varies marginally from its value for which the filling factor is exact. It turns out that c is the inverse of m , as the strength (magnetic charge) m with which the 2-dimensional system buffers a unit of extra flux is to create a quasi-hole of (electric) charge c so that $mc = 1$ flux unit. We shall call (11) the associated coupling matrix of m , and its being singular the auxiliary duality condition. A remark is imperative at this juncture. Strictly speaking, particles of different charges are distinguishable and the conventional notion of statistics, which concerns a system of particles being *indistinguishable*, does not apply *per se*. But as the representation theory of braid group does not require the charges to be identical, one can still talk about two *distinguishable* particles exchanging their locations. The so-called braid group “statistics” is topological in nature, for the configuration space for each particle is riddled with punctures and hence non-simply connected.

Now let us generalize the duality concept encapsulated in $mc = 1$ to the double-layered systems:

$$\left(\begin{array}{cc} m_1 & n \\ n & m_2 \end{array} \right) \left(\begin{array}{c} c_1 \\ c_2 \end{array} \right) = \left(\begin{array}{c} 1 \\ 1 \end{array} \right). \quad (12)$$

In other words, we have replaced m by the coupling matrix (7) and c by the charge vector (c_1, c_2) . In retrospect, the strength m for the single-layered FQHE should be considered as a 1×1 matrix and c a 1×1 vector. The duality condition determines

the values of the charges c_1, c_2 . The meaning of (12) has been discussed in ref. [12] by counting the total flux that each electron sees. For electrons of layer 1, it is $\Phi^{[1]} \equiv m_1(N^{[1]} - 1) + nN^{[2]}$, while every electron of layer 2 sees a total of $\Phi^{[2]} \equiv m_2(N^{[2]} - 1) + nN^{[1]}$ units of flux. The crucial point is that at the large scale limit, the two layers are in equilibrium and therefore $\Phi^{[1]} = \Phi^{[2]}$, which is

$$(N^{[1]} - 1)m_1 + N^{[2]}n = N^{[1]}n + (N^{[2]} - 1)m_2. \quad (13)$$

Now, quasi-holes behave much like electrons, except they are fractionally charged. Therefore, they also engage themselves in the puncture phase. So, in addition to $\Phi^{[1]}$ that each electron of tag [1] sees, it also senses the flux carried by the quasi-holes: $m_1c_1N_1 + nc_2N_2 = N_1$. Similarly, each electron of tag [2] senses an additional $m_2c_2N_2 + nc_1N_1 = N_2$ units of flux. Here, N_1 and N_2 denote the numbers of quasi-holes in layer 1 and layer 2 respectively. Since both layers are immersed in the same background magnetic field, small variation in the field strength should result in the same number of extra fluxes. In other words, $N_1 = N_2$. Therefore, we get the following two equations: $m_1c_1 + nc_2 = 1$ and $m_2c_2 + nc_1 = 1$. These equations are none other than the generalized duality condition (12). Upon solving the equations, we find $c_1 = (m_2 - n)/(m_1m_2 - n^2)$ and $c_2 = (m_1 - n)/(m_1m_2 - n^2)$. Having determined the values of the charges of the quasi-holes, we now calculate the filling factor of the double-layered FQHE. The filling factor of the quantum Hall effect is given by the total number of electrons divided by the total number of flux. In the single-layered case, the total number of flux that an electron sees is $(N - 1)m$. Hence, in the limit $N \rightarrow \infty$, we have $\frac{N}{(N-1)m} \rightarrow \frac{1}{m}$. Similarly, for the double-layered FQHE, using (13) we get $(N^{[1]} + N^{[2]})/(N^{[1]}m_1 + N^{[2]}n) \rightarrow (m_1 + m_2 - 2n)/(m_1m_2 - n^2)$ in the limit $N^{[1]} \rightarrow \infty$ and $N^{[2]} \rightarrow \infty$. With the physical picture offered by the path integral representation of the braid group, we can proceed to write down the many-quasi-hole wavefunction of the bilayered system:

$$\begin{aligned} \psi &\equiv \prod_{a < b} (u_a^{[1]} - u_b^{[1]})^{m_1c_1^2} \prod_{a, b} (u_a^{[1]} - u_b^{[2]})^{nc_1c_2} \prod_{a < b} (u_a^{[2]} - u_b^{[2]})^{m_2c_2^2} \\ &\times \prod (w_a^{[1]} - u_b^{[1]})^{m_1c_1} \prod (w_a^{[2]} - u_b^{[2]})^{m_2c_2} \prod (w_a^{[1]} - u_b^{[2]})^{nc_2} \prod (w_a^{[2]} - u_b^{[1]})^{nc_1} \\ &\times \exp\left(-\frac{1}{4\ell^2c_1} \sum_a^{N_1} |u_a^{[1]}|^2 - \frac{1}{4\ell^2c_2} \sum_a^{N_2} |u_a^{[2]}|^2\right) \psi_{m_1, m_2, n} \end{aligned} \quad (14)$$

This wavefunction is an exact solution of the bilayered version of the ground state equations: $(\mathcal{D}_a^{[i]} + \frac{c^{[i]}B}{4}\overline{u}_a^{[i]}) \psi = 0$ where

$$\mathcal{D}_a^{[1]} \equiv \partial_{u_a^{[1]}} - m_1 \sum_b^{N^{[1]}} \frac{c_1 \times 1}{u_a^{[1]} - w_b^{[1]}} - n \sum_b^{N^{[2]}} \frac{c_1 \times 1}{u_a^{[1]} - w_b^{[2]}} - m_1 \sum_{b \neq a}^{N_1} \frac{c_1 \times c_1}{u_a^{[1]} - u_b^{[1]}} - n \sum_b^{N_2} \frac{c_1 \times c_2}{u_a^{[1]} - u_b^{[2]}}$$

$$\begin{aligned}
&= \partial_{u_a^{[1]}} - m_1 c_1 \sum_b^{N^{[1]}} \frac{1 \times 1}{u_a^{[1]} - w_b^{[1]}} - n c_1 \sum_b^{N^{[2]}} \frac{1 \times 1}{u_a^{[1]} - w_b^{[2]}} \\
&\quad - m_1 c_1^2 \sum_{b \neq a}^{N_1} \frac{1 \times 1}{u_a^{[1]} - u_b^{[1]}} - n c_1 c_2 \sum_b^{N_2} \frac{1 \times 1}{u_a^{[1]} - u_b^{[2]}},
\end{aligned} \tag{15}$$

and a similar expression for $\mathcal{D}_a^{[2]}$ with $1 \leftrightarrow 2$. The statistical parameters of the quasi-holes are easily readable from (14) and they are

$$\theta^{[11]}/\pi = m_1 c_1^2, \quad \theta^{[12]}/\pi = n c_1 c_2, \quad \theta^{[22]}/\pi = m_2 c_2^2. \tag{16}$$

These values were first obtained by Wilczek [13] using the adiabatic transport argument. We have checked that these values coincide with those calculated with the generalized Chern-Simons approach [14]. $\theta^{[12]}/\pi$ is the *mutual statistics* of the quasi-holes residing in different layers. The associated coupling matrix of the double-layered systems is

$$\begin{array}{c}
\begin{array}{cccc}
& e^{[1]} & e^{[2]} & qh^{[1]} & qh^{[2]} \\
e^{[1]} & \left(\begin{array}{cccc}
m_1 & n & m_1 c_1 & n c_2 \\
n & m_2 & n c_1 & m_2 c_2 \\
m_1 c_1 & n c_1 & m_1 c_1^2 & n c_1 c_2 \\
n c_2 & m_2 c_2 & n c_1 c_2 & m_2 c_2^2
\end{array} \right)
\end{array}
\end{array} \tag{17}$$

One can check that this 4×4 matrix is singular, as befits the auxiliary duality condition for double-layered FQHE.

The generalization of the discussion to multi-sheeted FQHE is immediate [12]. Suppose the number of layers is M . Then, the braid group dynamics gives rise to a $M \times M$ coupling matrix which is non-singular, and its associated $2M \times 2M$ matrix which is singular. Using these matrices and the duality, we can characterize the FQHE on multi-layered samples. In particular, one should get a $\nu = 5/7$ for the triple-layered electronic systems with coupling matrix

$$\begin{pmatrix} 3 & 1 & 0 \\ 1 & 3 & 1 \\ 0 & 1 & 3 \end{pmatrix}. \tag{18}$$

Furthermore, it seems that one can even consider the following coupling matrix for

3-dimensional FQHE:

$$\begin{pmatrix} m_1 & n & 0 & \cdots & \cdots & 0 \\ n & m_2 & n & \cdots & \cdots & \vdots \\ 0 & n & m_3 & \ddots & & \vdots \\ \vdots & \vdots & \ddots & \ddots & \ddots & \vdots \\ \vdots & \vdots & & \ddots & m_{M-1} & n \\ 0 & 0 & \cdots & \cdots & n & m_M \end{pmatrix} \quad (19)$$

The symmetric tridiagonal coupling matrix (19) is an interesting case because it corresponds to the nearest-neighbour inter-layer correlation which is more likely to be realized experimentally. Now, when the number of layers M is large, the electronic system is literally 3-dimensional! By juxtaposing narrow quantum wells alternately with barriers of appropriate width, one may create 3-dimensional FQHE in the laboratory. The multi-layered FQHE presented here through the looking glass of coupling matrix may be relevant to the theory of anyonic superconductivity. So far, most of the effective theories with a Chern-Simons piece in the Lagrangian seem to ignore the inter-plane correlation. It may well turn out that the correlation between the copper oxide planes conspires to circumvent the P and T violation [13]. This is an open question.

In summary, we want to highlight the main theme of this work. We see that the coupling matrix and the associated coupling matrix contain all the relevant data that one needs to characterize the FQHE in samples with arbitrary number of layers. With the help of duality and the auxiliary duality condition, we can determine the (fractional) charges of the quasi-hole excitations, the statistics and the filling factor of the Hall media. This set of tools comes from the representation theory of the braid group over a multi-sheeted surface [12].

I would like to thank C. H. Lai for encouragement and discussions, and C. L. Ho for a useful conversation on “duality”.

Note added: After this paper was submitted for publication, I learned that Ezawa and Iwazaki had also independently considered the concept of mutual statistics (which they called relative statistics) in [15].

References

- [1] R. B. Laughlin, Phys. Rev. Lett. **50**, 1395 (1983).
- [2] Y. W. Suen, L. W. Engel, M. B. Santos, M. Shayegan and D. C. Tsui, Phys. Rev. Lett. **68**, 1379 (1992).
- [3] J. P. Eisenstein, G. S. Boebinger, L. N. Pfeiffer, K. W. West and Song He, Phys. Rev. Lett. **68**, 1383 (1992).
- [4] D. Yoshioka, A. MacDonald and S. Girvin, Phys. Rev. **B39**, 1932 (1988).
- [5] B. I. Halperin, Helv. Phys. Acta **56** 75, (1983).
- [6] C. Ting and C. H. Lai, Mod. Phys. Lett. **B5** 1293, (1991).
- [7] C. Ting and C. H. Lai, *Spinning Braid Group Representation and the Fractional Quantum Hall Effect*, NUS preprint NUS/HEP/92011, (Jan, 1992) (hep-th@xxx/ 9202024).
- [8] C. H. Lai and C. Ting, Phys. Lett. **B265**, 341 (1991).
- [9] Y. S. Wu, Phys. Rev. Lett. **52**, 2103 (1984).
- [10] E. Verlinde, *A note on braid statistics and the non-abelian Aharonov-Bohm effect*, preprint IASSNS-HEP-90/60, Jan 1991.
- [11] V. Knizhnik and A. Zamolodchikov, Nucl. Phys. **B247**, 83 (1984).
- [12] C. Ting, *Mutual statistics, braid group, and the fractional quantum Hall effect*, preprint NUS/HEP/920502, (May, 1992).
- [13] F. Wilczek, *Disassembling Anyons*, preprint IASSNS-HEP-91/70, Jan 1992.
- [14] A. Zee, in the *Proceedings of the TCSUH Workshop on Physics and Mathematics of Anyons*, edited by S. S. Chern, C. W. Chu and C. S. Ting, Int. J. Mod. Phys. **B5**, 1629 (1991); J. Fröhlich and A. Zee, Nucl. Phys. **B364**, 517 (1991).
- [15] Z. F. Ezawa and A. Iwazaki, *Chern-Simons Gauge Theory for Double-Layer Electron System*, preprint TU-402, Nisho-18 (May, 1992).